




## Chapter 2 An Introduction to Organic Compounds Functions, Nomenclature, Physical Properties, and Conformations Adapted from Profs. Turro & Breslow, Columbia University and Prof. Irene Lee, Case Western Reserve University














## Nomenclature of Alcohols

- In an alcohol, the OH is a functional group
- A functional group is the center of reactivity in a molecule
- Determine the parent hydrocarbon containing the functional group

CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>B

3-butoxy-1-propanol or 
3-butoxypropan-1-ol

1 2 3 4 3 2 1 CH3
HB 2H 2B B CICH2CH2CH3 5 14 3 2
3-bromo-1-propanol OH CH3 OH
4-chloro-2-butanol 4.4-dilmethyl-2-p

3. When there is both a functional group suffix and a substituent, the functional group suffix gets the lowest number


CH<sub>3</sub>CHCHCH<sub>2</sub>CH<sub>3</sub>
CI OH


2-chloro-3-pentanol not not -4-chloro-3-pentanol 6-methyl-4-heptanol


If there is more than one substituent, the substituents are cited in alphabetical order


CH<sub>2</sub>CH<sub>3</sub>


CH<sub>2</sub>CH<sub>3</sub>














Other Common Functional Groups	
Class	General Formula
Aldehydes	R-C-H
Ketones	Q R-C-R'
Carboxylic Acids	О R-C-OH
Esters	Q R-C-OR'
Amides	O R" R-C-N R'

Attractive Forces	
Ionic bonds	
Covalent bonds	
Hydrogen bonds	
Dipole-dipole interaction	
lon-dipole	
van der Waals force Dispersion Forces	
The greater the attractive intermolectar forces between molecules, the higher is the boiling point of the compound, eg. water.	

